

10018604



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

Publication number:

**0 271 988  
A2**

12

# EUROPEAN PATENT APPLICATION

21 Application number: 87309853.7

51 Int. Cl. 4: **C12N 15/00** , **C12N 5/00** ,  
**A01H 1/00**

22 Date of filing: 06.11.87

The applicant has filed a statement in accordance with Rule 28 (4) EPC (issue of a sample only to an expert). Accession number(s) of the deposit(s): NCIB-12352, NCIB-12373, NCIB-12568.

30 Priority: 11.11.86 GB 8626879

43 Date of publication of application:  
22.06.88 Bulletin 88/25

84 Designated Contracting States:  
**AT BE CH DE ES FR GB GR IT LI LU NL SE**

71 Applicant: **IMPERIAL CHEMICAL INDUSTRIES PLC**  
**Imperial Chemical House Millbank**  
**London SW1P 3JF(GB)**

72 Inventor: **Bridges, Ian George**  
**Greentrees**  
**Manley Lane Cheshire(GB)**  
Inventor: **Schuch, Wolfgang Walter**  
**20 Belmonte Road Overton**  
**Frodsham Cheshire(GB)**  
Inventor: **Grierson, Donald**  
**6 Tyler Court**  
**Shepshed Leicestershire(GB)**

74 Representative: **Roberts, Timothy Wace et al**  
**Imperial Chemical Industries PLC Legal**  
**Department : Patents PO Box 6 Bessemer**  
**Road**  
**Welwyn Garden City Herts, AL7 1HD(GB)**

54 **Anti-sense regulation of plant gene expression.**

57 Recombinant DNA comprises promoter and terminator base sequences respectively upstream and downstream of an inverted base sequence complementary to a substantial sequence of bases in polygalacturonase mRNA. The antisense mRNA produced thereby delays softening of fruit, in particular tomatoes.

EP 0 271 988 A2

DNA

This invention relates to DNA: and to vectors and plant cells containing such DNA, and to plants composed of such cells. In one aspect, the invention relates to flowering and fruiting plants, for example tomatoes, and to such plants having usefully altered ripening properties.

It has been shown that tomato ripening is a process involving the activation of specific genes (D. Grierson, 1985, CRC Critical Reviews in Plant Sciences 3, 113-132). The features of many of these genes have not been defined. However, one such gene has been shown to produce polygalacturonase (PG), the enzyme primarily responsible for degrading the cell wall. The synthesis of PG begins in tomatoes during early stages of ripening, and reaches a maximum at the soft red stage. This increase is paralleled closely by an increase in PG mRNA (Grierson et al, 1985, Planta 163, pp. 263-271).

Various investigations, including enzyme analysis, indicate that the PG gene, in tomatoes, is only expressed in ripening fruit (Maunder et al, 1986, in press) and in flower abscission zones.

A hitherto unpublished paper of which the present inventors are authors (Grierson et al, 1986, Nucleic Acid Reviews, 14 p 8595-8603) describes the preparation of cDNA from PG mRNA, and discloses in particular pTOM6, a plasmid containing substantially all (all but the first 20 bases) of the cDNA sequence complementary to the mRNA that is generated in the ripening tomato and that is translated into PG.

We now propose to regulate the expression of the PG gene in flowering and fruiting plants (such as tomatoes) by generating antisense RNA to PG sequences in such fruit. Such RNA hinders the expression of the PG mRNA, probably by forming therewith a double-stranded complex.

It has previously been proposed to regulate gene expression in both prokaryotes and eukaryotes by generating antisense RNA therein; and such proposals have included higher plants (Ecker et al, 1986, PNAS 83, pp. 5372-5376). However, we are aware of no prior proposals for using antisense RNA corresponding to a natural plant gene to control plant biochemistry or development.

According to the present invention we provide recombinant DNA comprising an upstream promoter base sequence, a base sequence for transcription into mRNA under control of said upstream promoter base sequence, and a downstream transcription terminator base sequence, characterised in that the base sequence for transcription comprises an inverted sequence of bases complementary to a substantial run of bases in mRNAs encoding fruit softening enzymes.

Examples of fruit softening enzymes are polygalacturonase and pectin methylesterase.

We further provide vectors containing such DNA; method of making such vectors by cloning of the desired base sequence for transcription in the appropriate orientation into an existing vector containing the desired promoter and terminator base sequences; plant cells containing such DNA, as well as plants (in particular tomatoes) constituted of such plant cells.

DNA according to the invention preferably comprises a base sequence for transcription at least 50 bases in length. There is no theoretical upper limit to the base sequence - it may be as long as the relevant mRNA produced by the cell - but for convenience it will generally be found suitable to use sequences between 100 and 1000 bases in length.

The invention will be further described with reference to the drawings, in which :

Figure 1 gives the base sequence for the polygalacturonase cDNA clone pTOM6;

Figure 2 shows schematically the fragments of pTOM6 and of the tomato polygalacturonase gene clone gTOM23 used in the following Examples and Experiments;

Figure 3 gives the base sequence for the pectin esterase cDNA clone pPE1.

A very convenient source of DNA for use as the base sequence for transcription is provided by DNA that gives rise to polygalacturonase mRNA.

The required antisense DNA can be obtained by cutting with restriction enzymes an appropriate sequence of such (mRNA-polygalacturonase producing) DNA; and then cloning the cut DNA into a vector containing upstream promoter and downstream terminator sequences, the cloning being so carried out that the cut DNA sequence is inverted with respect to its orientation in the strand from which it was cut.

In the new vector, the strand that was formerly the template strand becomes the coding strand, and vice versa. The new vector will thus produce RNA in a base sequence which is complementary to the sequence of polygalacturonase mRNA. Thus, the two RNA strands are complementary not only in their base sequence but also in their orientations (5' to 3').

As source of the polygalacturonase DNA base sequence for transcription, it is convenient to use a cDNA clone such as pTOM6. The base sequence of pTOM6 is set out in Figure 1. A cDNA clone such as pTOM6 may be obtained from the mRNA of ripening tomatoes by the method described by Slater et al, Plant Molecular Biology 5, 137-147. In this way may be obtained sequences coding for the whole, or

substantially the whole, of the mRNA that is translated into PG. Suitable lengths of the cDNA so obtained may be cut out for use by means of restriction enzymes.

An alternative source of DNA for the base sequence for transcription is the PG gene. It differs from that in the cDNA of, eg. pTOM6, in that introns are present. The introns are not transcribed into mRNA (or, if so transcribed, are subsequently cut out). The major part of the PG gene has been deposited as gTOM23 with the National Collections of Industrial and Marine Bacteria, Aberdeen, under Accession No. 12373. When using the PG gene as the source of the base sequence for transcription it is preferred to use primarily exon regions.

A further way of obtaining a suitable DNA base sequence for transcription is to synthesise it ab initio from the appropriate bases, for example using Figure 1 as a guide.

Recombinant DNA and vectors according to the present invention may be made as follows. A suitable vector containing the desired base sequence for transcription (for example pTOM6) is treated with a restriction enzyme to cut the sequence out. The DNA strand so obtained is cloned (in reverse orientation) into a second vector containing the desired promoter sequence (for example CAMV 35S or the PG gene promoter sequence) and the desired terminator sequence (for example the 3' end of the nopaline synthase gene, the nos 3' end).

It is often preferred, in applying the invention, to use the promoter of the PG gene. Use of this promoter, at least in tomatoes, has the advantage that the production of antisense RNA is under the control of the same system that controls PG mRNA. Thus whatever factors tend to produce the latter will tend at the same time to produce the former to interfere with it.

Vectors according to the invention may be used to transform plants as desired, to make plants according to the invention. Dicotyledonous plants, such as the tomato, may be transformed by Ti plasmid technology, for example as described by Bevan (1984) *Nucleic Acids Research*, 12, 87118721.

Such transformed plants may be reproduced sexually, or by cell culture.

The degree of production of antisense RNA in the plant cells can be controlled by suitable choice of promoter sequences, or by selecting the number of copies, or the site of integration, of the DNA sequences according to the invention that are introduced into the plant genome. In this way, for example, it may prove possible to delay softening of tomatoes for a greater or lesser period after ripening.

The following Examples and Experiments illustrate the invention and how to carry it out. Examples relate to the manufacture of vectors according to the invention: Experiments relate to preparation of starting materials. All cloning procedures are performed under standard conditions as described by Maniatis et al (1982) "Molecular Cloning", Cold Spring Harbor Laboratory. Vectors for which an NCIB Accession number are given have been deposited at the National Collections of Industrial and Marine Bacteria, Torry Research Station, PO Box 31, 135 Abbey Road, Aberdeen AB9 8DG, Scotland.

## EXPERIMENT 1

### Construction of the plasmid pPH1

#### A. Isolation of the nos 3' end

10 ug of pWRECK2-CAMV-CAT-1 (NCIB Accession No. 12352) is digested with PvuI in order to linearise the DNA, under conditions recommended by the manufacturer. The completeness of digestion is analysed by running an aliquot of the reaction of 0.8% agarose gels. The reaction is stopped by extraction with phenol/chloroform. DNA is precipitated with ethanol and dried under vacuum. The cohesive ends are removed by incubation of the linearised DNA with T4 polymerase at 37°C for 30 minutes. The enzyme is inactivated by incubation at 65°C for 15 minutes. The reaction volume is increased by the addition of HindIII buffer and HindIII enzyme is added. The reaction is carried out for 2 hours at 37°C. The 250 bp PvuI/HindIII fragment is isolated from agarose gels by electroelution. DNA is phenol/chloroform extracted, precipitated with ethanol and resuspended in water.

## B. Linearisation of pUC18.

2 ug of pUC18 (plasmid commercially available eg. from Amersham) DNA is digested with Sph1 under conditions recommended by the manufacturer. The reaction is stopped by extraction with phenol/chloroform. Cohesive ends are removed by treatment with T4 polymerase for 30 minutes at 37°C. The buffer volume is increased, and HindIII is added. The mixture is incubated for 2 hours at 37°C. The reaction is stopped by extraction with phenol/chloroform. DNA is precipitated with ethanol and resuspended in water at 100 ng/ml.

## C. Cloning of nos. 3' end into pUC18 to give pNOS1.

1 ul of pUC18 prepared under (B) is ligated with 100 ng of nos 3' end prepared under (A) in a total of 15 ul in the presence of T4 ligase. Incubation is carried out for 24 hours at 16°C. An aliquot of the ligation is transformed into competent TG2 cells. An aliquot of the transformation mix is plated onto ampicillin and Xga1 containing plates. Clear plaques are picked, and the DNA examined by restriction analysis. Molecules containing the nos 3' end are characterised by the presence of a 260 base pair HindIII/BamHI fragment. These plasmids are called pNOS1.

## D. Preparation of the CaMV 35S promoter

The CaMV promoter is obtained by digestion of pWRECK2-CAMV-CAT-1 (NCIB Accession No. 12352) with Scal for 2 hours at 37°C. An aliquot of the restriction digest is analysed by electrophoresis on agarose gels. The reaction is stopped by extraction with phenol/chloroform. After ethanol precipitation and resuspension in water, the DNA is cut with Hph1 for 2 hours at 37°C. The cohesive ends are removed by treatment with T4 polymerase under standard conditions. The CaMV promoter 629 base pair fragment is isolated by agarose gel electrophoresis and subsequent electroelution.

## E. Linearisation of pNOS1

2 ug of pNOS1 is cut with Sst1 at 37°C for 2 hours. After completion of the reaction, T4 polymerase is added in order to remove the cohesive ends. The reaction is stopped by extraction with phenol/chloroform. Then DNA is precipitated with ethanol and resuspended in water at 100 ng/ul.

## F. Cloning of CaMV 35S promoter into pNOS1

pNOS1 prepared as under (E) is ligated to CaMV 35 promoter fragment prepared under (D) under standard conditions using T4 ligase. The reaction is carried out for 24 hours at 16°C. An aliquot of the ligation mixture is transformed into competent TG2 cells, and plated onto ampicillin containing Xga1 plates. DNA is isolated from transformants and analysed by restriction with NcoI and HindIII. Molecules containing the CaMV 35S promoter in the correct orientation are characterised by the presence of a 920 base pair fragment. These plasmids are called pPH1.

## EXPERIMENT 2 : Preparation of Plasmid pCB1

## A. Isolation of a PG promoter fragment

Genomic clones are isolated from a partial Sau3A library of Ailsa Craig tomato DNA cloned into EMBL3 (Bird et al, in preparation). PG clones are isolated from the genomic library by screening with both the complete pTOM6 cDNA insert, and the isolated 5' Pst1/HindIII fragment from pTOM6 (Grierson et al, NAR 1986). Several overlapping clones are isolated and the transcription start site of the PG gene located by S1 mapping experiments (Bird et al in preparation 1987). The PG promoter can be located on a 1.6 Kb HindIII fragment which also contains part of the PG coding information.

## B. Insertion of a SpeI site into the PG promoter

In order to be able to manipulate the PG promoter sequence conveniently (ie. the DNA 5' to the transcription start) a SpeI site is introduced by site directed mutagenesis using standard protocols. The  
 5 HindIII fragment is isolated from genomic clone gTOM23 (NCIB Accession No. 12373), and cloned into the HindIII site of (commercially available vector) M13 mp19. After annealing with a suitable mismatch primer and extension using DNA polymerase, the mixture is transformed into competent TG2 cells. Phages are plated and duplicated filters were prepared for hybridisation to the labelled mismatch primer. Putative  
 10 clones are identified by hybridisation under increasingly stringent conditions, isolated and the generation of the SpeI site is determined by direct DNA sequence analysis. The promoter fragment is isolated from one isolate by restriction with SpeI and HindIII. This fragment is then cloned into pUC19 (commercially available plasmid) cut with HindIII and XbaI. The promoter fragment is then transferred into Bin19 (Bevan, Nucleic Acid Research, 1984, 12, 87118721) cut with BamHI and HindIII. This plasmid is called pCB1.

15

## EXPERIMENT 3 : Preparation of Plasmid pPH2

### A. Isolation of the PG promoter fragment from pCB1

20 5 ug of pCB1 (prepared as in Experiment 2) is cut with HindIII for 2 hours at 37°C. The mixture is phenol/chloroform extracted and DNA precipitated with ethanol. After re-suspension in water the cohesive ends are filled in using DNA polymerase under standard conditions at room temperature for 15 minutes. The polymerase is inactivated by heating to 65°C for 15 minutes. The DNA is then treated with BamHI for  
 25 2 hours at 37°C. The PG promoter fragment is then by electroelution isolated by agarose gel electrophoresis as a HindIII/BamHI 1.45 Kb fragment.

### B. Preparation of pPH1 for insertion of the PG promoter fragment

30 5 ug of pPH1 (prepared as in Experiment 1) is cut with NcoI for 2 hours at 37°C under standard conditions. The DNA is purified by phenol/chloroform extraction. The cohesive ends are filled in using DNA polymerase I Klenov fragment for 15 minutes at room temperature. The volume is increased and BamHI added. The mixture is incubated for 2 hours at 37°C. The mixture is then separated on agarose gels, and the large fragment of approximately 3 Kb isolated by electroelution.

35

### C. Cloning of the PG promoter into the large fragment from pPH1.

40 10 ug of pPH1 prepared as in Experiment 1 is ligated with the PG promoter fragment as prepared in A under standard conditions for 24 hours at 16°C. An aliquot of the ligation mixture is used to transform competent TG2 cells. Aliquots of the transformation mixture are plated onto L plates containing ampicillin and Xga1. Colonies are picked and examined for the presence of the PG promoter DNA by electrophoresis to detect an increase in the size of the vector and by direct DNA sequence determination. This plasmid is called pPH2.

45

## EXAMPLE 1 : Preparation of Plasmids pJR10 and pJR11

### A. Preparation of the antisense DNA

50

A 730 base pair HinfI fragment covering the 5' untranslated region, the putative leader sequence, and a substantial portion of the PG coding region is isolated from pTOM6. 5 ug pTOM6 is restricted with HindI for 2 hours at 37°C. The reaction is stopped by extraction with phenol/chloroform, and ethanol precipitated. Cohesive ends are removed by treatment with T4 polymerase under standard conditions. DNA is purified by  
 55 phenol/chloroform extraction and ethanol precipitation.

## B. Cloning of the antisense DNA into pPH1

The DNA fragment as isolated in Example 1.A is cloned into the Sma1 site of pJR1. pPH1 DNA is restricted with Sma1 under standard conditions for 2 hours at 37°C. After incubation for 2 hours bacterial alkaline phosphatase is added in order to prevent self-ligation of pPH1 during subsequent cloning steps. The reaction is stopped by extraction with phenol/chloroform. DNA is precipitated and resuspended in water. An aliquot of the DNA (pPH1) is ligated under standard conditions to Sma1 cut pPH1. Aliquots of the ligation mixture are transformed into competent TG2 cells, and plated onto ampicillin containing plates. Recombinants are analysed by restriction digestion with HindIII. Both antisense (pPH10) and "sense" (pPH11) constructs are isolated.

## EXAMPLE 2 : Preparation of Plasmids pPH20 and pPH21

### 15 A. Cloning of the antisense DNA into pPH2

DNA as isolated in Example 1A is cloned into pPH2 cut with HincII. The plasmid containing the PG sequence in the antisense orientation is called pPH20, the plasmid containing the PG DNA in the sense orientation is called pPH21.

## EXAMPLE 3 : Transformation of Tomato Plants

Transformation of tomato plants is achieved using a modification of the leaf disc transformation protocol published by Bevan et al, EMBO Journal 4, 1921-1926, 1985. Transformed tomato plants are analysed for the presence of the antisense constructs by Southern hybridisation to genomic DNA. Expression of sense and antisense RNA is monitored by dot blot and Northern hybridisations. Firmness of the fruit is investigated using general physiological methods. The presence of antisense PG constructs in the cells of the ripening tomatoes is associated with firmness in the tomato being maintained for a longer period.

There follows a second series of Examples and Experiments which have been carried out according to the same general scheme.

## EXPERIMENT 11

### 35 Contruction of the plasmid pJR1

#### A. Isolation of the nos. 3' end

This was carried out according to the method of Experiment 1 above.

#### B. Removal of the CaMV 3' end from pDH51

2-μg of pDH51 (Pietrzak et al, (1986) Nucleic Acids Research 14, 5857-5868) was digested with SphI at 37°C for 2 hours under standard conditions. The reaction was stopped by extraction with phenol/chloroform. DNA was precipitated with ethanol and resuspended in water.

Cohesive ends were removed by treatment with T4 polymerase for 30 minutes at 37°C. The buffer volume was increased, and HindIII was added. The mixture was incubated for 2 hours at 37°C. The resulting 3.2 Kb fragment was isolated after gel electrophoresis on agarose gels by electroelution. The DNA was extracted with phenol and chloroform, precipitated with ethanol and resuspended in water.

### C. Cloning of nos 3' end into pDH51 to give pJR1

1  $\mu$ l of pDH51 prepared under (B) was ligated with 100 ng of nos 3' end prepared under (A) in a total of 15  $\mu$ l in the presence of T4 ligase. Incubation was carried out for 24 hours in 16°C. An aliquot of the ligation was transformed into competent TG2 cells. An aliquot of the transformation mix was plated onto ampicillin and Xga1 containing L-plates. White colonies were picked, and the DNA examined by restriction analysis. Molecules containing the nos 3' end were characterised by the presence of a 260 base pair HindIII - BamHI fragment. These plasmids were called pJR1.

10

## EXPERIMENT 12

Construction of plasmids pDHC1 and pDHC4.

### 15 A. Isolation of a 730 bp HinfI fragment from pTOM6

5  $\mu$ g pTOM6 (NCIB accession No. 12351) was treated with HinfI for 2 hours at 37°C under standard conditions. The 730 bp HinfI fragment (Figure 3) was isolated after separation on agarose gels. The cohesive ends of this fragment were filled in with DNA polymerase Klenov fragment A. The DNA was phenol extracted and ethanol precipitated.

20

### B. Linearisation of pDH51

25 1  $\mu$ g pDH51 was treated with SmaI for 2 hours at 37°C under standard conditions. The reaction was stopped by phenol extraction. The linearised vector was then precipitated with ethanol, washed and resuspended in water.

25

### 30 C. Cloning of the pTOM6 HinfI fragment into pDH51

The isolated HinfI fragment from pTOM6 (A) and the linearised vector (B) were ligated overnight under standard conditions. The ligation mix was used to transform competent TG2 cells. The transformation mix was plated onto ampicillin-containing plates. Clones were selected, DNA isolated and analysed by digestion with BamHI and HindIII restriction enzymes. Plasmids were identified, and were named pDHC1 and pDHC4. pDHC1 contains the HinfI fragment in the antisense orientation; pDHC4 contains the HinfI fragment in the sense orientation.

35

## 40 EXPERIMENT 13

Construction of plasmid pCB1.

### A. Isolation of a PG promoter fragment

45

This was carried out as described in Experiment 2A above.

### B. Insertion of a Spe1 site into the PG promoter fragment.

50

This was carried out as described in Experiment 2B above. The resulting plasmid was called pCB1.

## EXPERIMENT 14

55

Construction of plasmid pJR2

## A. Isolation of the PG promoter fragment from pCB1

This was carried out as described in Experiment 3A.

5

## B. Preparation of pJR1 for insertion of the PG promoter fragment.

5µg of pJR1 (constructed in Experiment 11) was cut with NcoI for 2 hours at 37°C under standard conditions. The DNA was purified by extraction with phenol and chloroform. The cohesive ends were filled in using DNA polymerase I Klenow fragment A for 15 minutes at room temperature. The volume was increased and BamHI added. The mixture was incubated for 2 hours at 37°C. The mixture was then fractionated on agarose gels, and the large fragment of approximately 3kb isolated by electroelution.

## C. Cloning of the PG promoter into the large fragment from pJR1.

pJR1 prepared as in B above was ligated with the PG promoter fragment prepared in A under standard conditions for 24 hours at 16°C. An aliquot of the ligation mixture was used to transform competent TG2 cells. Aliquots of the transformation mixture were plated onto L plates containing ampicillin and Xga1. Colonies were picked and examined for the presence of the PG promoter DNA by electrophoresis on agarose gels in order to detect an increase in the size of the vector and by direct DNA sequence determination. Plasmids containing the PG promoter were called pJR2.

## CONSTRUCTION OF ANTISENSE AND SENSE PG VECTORS

A series of antisense and sense (control) vectors containing different portions of the PG cDNA and PG gene were constructed for use in regenerating transgenic plants. The vectors produced are summarised in Table 1. The vectors constructed are based on pJR1 and pJR2. DNA fragments have been inserted into these vectors both into the antisense (A) and sense (B) orientations. Expression cassettes contained in these vectors were then transferred to Bin19 (Bevan (1984) Nucleic Acids Research, 12, 8711-8721) for transformation of tomato plants.

35

TABLE 1

40	Vectors based on	Name of antisense vector	Name of sense vector	PG fragment (see Figure 2)
45	pJR1	pJR16A pJR36A pJR56A pJR76A	pJR16S pJR36S pJR56S pJR76S	740 bp HinfI fragment a fragment b fragment c
50	pJR2	pJR26A pJR46A pJR66A pJR86A	pJR26S pJR46S pJR66S pJR86S	740 bp HinfI fragment a fragment b fragment c
55				



## EXAMPLE 10

## Construction of PG antisense vectors pJR16A and pJR16S

## 5 A. Isolation of a 740 bp PG antisense fragment

5 g pDHC1 was cut with KpnI and PstI at 37°C for 2 hours under standard conditions. The 740 bp KpnI - PstI fragment was isolated after agarose gel electrophoresis by electroelution. The fragment was extracted with phenol and chloroform and ethanol precipitated. The fragment was then resuspended in 10µl TE.

10

## B. Preparation of pJR1

15 1µg pJR1 (from Experiment 14) was cut with KpnI and PstI at 37°C for 2 hours. The reaction was stopped by extraction with phenol and chloroform. The DNA was precipitated with ethanol, washed and dried. The vector was resuspended in 20µl TE.

## C. Ligation of the PG antisense fragment and pJR1

20

The products of (A) and (B) above were ligated at 16°C for 24 hours under standard conditions. The ligation was used to transform competent TG2 cells and the mixture was plated onto ampicillin-containing plates to select transformed cells. Single colonies were grown up to prepare plasmid DNA. The DNA was analysed for the presence of 500 bp HindIII fragment. A clone containing this fragment was identified and called pJR16A.

25

## pJR16S

## 30 D. Isolation of a 740 bp PG sense fragment

5 g pDHC4 was cut with KpnI and PstI at 37°C for 2 hours under standard conditions. The 740 bp fragment produced was isolated after agarose gel electrophoresis by electroelution. The fragment was extracted with phenol and chloroform and precipitated with ethanol. The fragment was then suspended in 10µl TE.

35

## F. Ligation of the PG sense fragment to pJR1

40 The products of (D) were ligated at 16°C for 24 hours under standard conditions. The ligation mix was used to transform competent TG2 cells and the mixture was plated onto ampicillin containing plates. Single colonies were grown up to prepare plasmid DNA. The DNA was analysed for the presence of a 900 bp HindIII fragment. A suitable clone was identified and called pJR16S.

45

## EXAMPLE 11

## Transfer of pJR16A and pJR16S to Bin 19

## 50 A. Isolation of the 1600 bp expression cassettes

pJR16A and pJR16S were cut with EcoRI at 37°C for 2 hours under standard conditions. An aliquot of the reaction mixture was separated by agarose gel electrophoresis to check that the reaction had gone to completion. It was then heated to 65°C for 15 minutes in order to inactivate the enzyme. The DNA was then cut partially with a small amount of HindIII in order to give all the possible EcoRI/HindIII partial digestion

55

fragments. The EcoRI - HindIII fragment of approximately 1600 bp consisting of the 35S CaMV promoter, the PG insert sequences and Nos 3' end (expression cassette) was isolated after agarose gel electrophoresis by electroelution. The fragment was extracted with phenol and chloroform, and precipitated with ethanol. The fragment was washed, dried and resuspended in 10µl TE.

5

#### B. Preparation of Bin19 for cloning

5 g Bin19 DNA was cut with EcoRI and HindIII at 37°C for 2 hours under standard conditions. The reaction was stopped by phenol and chloroform extraction, and DNA was precipitated with ethanol. The vector prepared in this fashion was resuspended in 20µl.

10

#### C. Ligation of Bin19 to PG expression cassettes

15

The products of (A) and (B) were set up for ligation. Aliquots of the PG antisense and sense cassettes were ligated to Bin19 at 16°C for 24 hours under standard conditions. The ligation mixes were used to transform competent TG2 cells which were plated on L agar containing Kanamycin and Xga1. Recombinant colonies were identified by their white colour. A number of these were picked from each ligation reaction and used to prepare plasmid DNA. The DNA was analysed for the relevant restriction pattern by cutting with EcoRI and HindIII.

20

#### EXAMPLE 12

25

#### Construction of PG antisense vectors pJR26A and pJR26S

##### pJR26A

#### 30 A. Isolation of a 740 bp PG antisense fragment

5 g pDHC1 were cut with KpnI at 37°C for 2 hours under standard conditions. The cohesive ends of the molecule were filled in with T4 DNA polymerase under standard conditions. The reaction was stopped by heating at 65°C for 15 minutes. The DNA was then also cut with PstI at 37°C for 2 hours under standard conditions. The resulting 740 bp KpnI (blunt) - PstI fragment was isolated after agarose gel electrophoresis by electroelution. The fragment was extracted with phenol and chloroform, and precipitated with ethanol. The fragment was then resuspended in 10µl TE.

35

#### 40 B. Preparation of pJR2

1 g pJR2 (from Experiment 14) was cut with HincII and PstI at 37°C for 2 hours under standard conditions. The reaction was terminated by extraction with phenol and chloroform and precipitated with ethanol. The purified vector was resuspended in 20µl TE.

45

#### C. Ligation of PG fragments to pJR2

The products of (A) and (B) above were ligated at 16°C for 24 hours under standard conditions. The ligation mix was used to transform competent TG2 cells. The transformation mix was plated onto ampicillin-containing plates and incubated at 37°C overnight. Transformed colonies were grown up and plasmid DNA was prepared for analysis. A clone was identified which contained a 2 Kb EcoRI - HindIII insert. This clone was called pJR26A.

50

55

##### pJR26S

#### D. Isolation of the 740 bp PG fragment

5  $\mu$ g pDHC4 were cut with KpnI at 37°C for 2 hours under standard conditions. The cohesive ends of the DNA were filled in with T4 DNA polymerase. The reaction was stopped by heating to 65°C for 15 minutes. The DNA was then also cut with PstI. The resulting 740 bp fragment was isolated after agarose gel electrophoresis by electroelution. The fragment was extracted with phenol and chloroform and precipitated with ethanol. It was then resuspended in 10  $\mu$ l TE.

#### 10 E. Ligation of the PG sense fragment to pJR2

The products of (B) and (D) above were ligated at 16°C for 24 hours under standard conditions. The ligation mix was used to transform competent TG2 cells, plated onto ampicillin containing plates and incubated at 37°C overnight. Transformed single colonies were grown up and plasmid DNA was prepared. 15 The DNA was analysed for the presence of a 1.6 Kb EcoRI - HindIII fragment. A clone was identified and called pJR26S.

### EXAMPLE 13

20

#### Transfer of pJR26A and pJR26S to Bin19

A procedure essentially the same as described above in Example 11 was used to subclone the 2.6 Kb EcoRI - HindIII partial fragments from pJR26A and pJR26S into Bin19 cut with EcoRI and HindIII. 25 Recombinants were identified by their white colour reaction after plating onto L-agar plates containing kanamycin and Xga1. Recombinants were characterised by restriction digestion with EcoRI and HindIII.

### EXAMPLE 14

30

#### Construction of vectors pJR36A, pJR36S (fragment a) and pJR46A and pJR46S (fragment b)

##### A. Isolation of fragments (a) and (b)

35 5  $\mu$ g pDHC4 (from Experiment 12) was cut with KpnI and BamHI at 37°C for 2 hours under standard conditions. The 500 bp fragment was isolated after agarose gel electrophoresis by electroelution, extracted with phenol, chloroform and resuspended in 20 l TE. The KpnI - BamHI fragment was then cut with HindIII. The cohesive ends of the fragment were filled with T4 DNA polymerase. The resulting fragments: a) 199 bp HindIII - KpnI (blunt ended) and b) 275 bp HindIII - BamHI (blunt ended) were isolated after agarose gel 40 electrophoresis by electroelution, extracted with phenol and chloroform, and resuspended in 10 l TE.

##### B. Preparation of pJR1

45 1  $\mu$ g pJR1 (from Experiment 11) was cut with SmaI at 37°C for 2 hours under standard conditions. The reaction was stopped by extraction with phenol and chloroform, and precipitated with ethanol. The vector was then resuspended in 20  $\mu$ l TE.

##### 50 C. Ligation of fragment (a) into pJR1

##### pJR36A and pJR36S

Fragment (a) from (A) above was ligated to SmaI cut pJR1 (from (B) above) at 16°C for 24 hours under 55 standard conditions. The ligation mixture was used to transform competent TG2 cells which were then plated onto ampicillin-containing plates. Transformed colonies were grown up and used for plasmid DNA

preparation. EcoRI/PstI double digests identified those clones containing fragment (a) inserts. The EcoRI - PstI inserts of these clones were isolated and subcloned into M13 mp8 which had been cut with EcoRI and PstI. DNA sequence analysis was carried out in order to ascertain the orientation of the insert (a). Clones obtained from this experiment were called pJR36A and pJR36S, according to the orientation of the insert.

#### D. Ligation of fragment (b) into pJR1

pJR56A and pJR56S

Fragment (b) from (A) above was ligated to SmaI cut pJR1, from (B) above, at 16°C for 24 hours under standard conditions. The ligation mixture was used to transform competent TG2 cells which were then plated onto ampicillin containing plates. Transformed colonies were grown up and used for plasmid DNA preparation. EcoRI/pstI double digests identified those clones containing fragment (b) inserts. The EcoRI - PstI inserts of these clones were isolated and subcloned into M13 mp8 which had been cut with EcoRI and PstI. DNA sequence analysis was carried out in order to ascertain the orientation of the insert (b). Clones obtained from this experiment were called pJR56A and pJR56S, according to the orientation of the insert.

### EXAMPLE 15

Transfer of pJR36A/S and pJR56A/S to Bin19

#### A. Preparation of expression cassettes containing fragments (a) and (b) in pJR1

5µg each of pJR36A, pJR36S, pJR56A and pJR56S were cut separately with EcoRI and HindIII at 37°C for 2 hours under standard conditions. The resulting four fragments containing (a) 930 bp and (b) 1000 bp were isolated separately after electrophoresis on agarose gels by electroelution. The fragments were extracted with phenol and chloroform, and precipitated with ethanol. The four fragments were then resuspended in 10µl TE.

#### B. Preparation of Bin19

Bin19 was cut with EcoRI and HindIII for 2 hours at 37°C under standard conditions. The reaction was stopped by addition of phenol and chloroform. After extraction the DNA was precipitated with ethanol, and resuspended in 20µl TE.

#### C. Ligation of the fragments to Bin19

The four EcoRI - HindIII fragments isolated in A were set up for separate ligation reactions using Bin19 prepared as described in B under standard conditions. The ligation mixtures were used to transform competent TG2 cells which were plated onto L agar containing kanamycin and Xga1. After incubation overnight, recombinant colonies were identified by their white colour. A number of the clones were picked from each separate ligation and were used to prepare DNA. The DNA's were analysed for the presence of a EcoRI - HindIII fragment of the appropriate size for the insertion of the expression cassettes to Bin19.

### EXAMPLE 16

Construction of pJR46A, pJR46S (fragment a) and pJR66A, pJR66S (fragment b)

#### A. Preparation of pJR2

1µg pJR2 (from Experiment 14) was cut with HincII at 37°C for 2 hours under standard conditions. The reaction was terminated by extraction with phenol and chloroform. The vector was precipitated with ethanol, washed and resuspended in 20µl TE.

pJR46A and pJR46S

#### B. Ligation of PG fragment (a) to pJR2

Fragment (a) from Example 14(A) above was ligated to HincII cut pJR2 from (A) above at 16°C for 24 hours under standard conditions. The ligation mixture was used to transform competent TG2 cells which were then plated onto ampicillin containing plates. Transformants were picked, grown up and used to prepare plasmid DNA. Plasmid DNA was cut with both EcoRI and PstI. DNA from clones which contained inserts were restricted with EcoRI and PstI. The EcoRI - PstI inserts were isolated after agarose gel electrophoresis by electroelution and subcloned into M13mp8 which had been cut with EcoRI and PstI. DNA sequence analysis was used to ascertain the orientation of the inserts (a). Clones were obtained from this experiment were called pJR46A and pJR46S, according to the orientation of the insert.

pJR66A and pJR66S

#### C. Ligation of PG fragment (b) to pJR2

Fragment (b) from Example 14(A) was ligated to HincII cut, from A above, separately at 16°C for 24 hours under standard conditions. The ligation mixture was used to transform competent TG2 cells which were then plated onto ampicillin containing plates. Transformants were picked, grown up and used to prepare plasmid DNA. Plasmid DNA was cut with both EcoRI and PstI. DNA from clones which contained inserts were restricted with EcoRI and PstI. The EcoRI - PstI inserts were isolated after agarose gel electrophoresis by electroelution and subcloned into M13 mp8 which had been cut with EcoRI and PstI. DNA sequence analysis was used to ascertain the orientation of the inserts (b). Clones were obtained from this experiment were called pJR66A and pJR66S, according to the orientation of the insert.

### EXAMPLE 17

Transfer of pJR46A, pJR46S, pJR66A and pJR66S to Bin19

#### A. Preparation of expression cassettes containing fragments (a) and (b) in pJR2

5 g of each of pJR46A, pJR46S, pJR66A and pJR66S were cut separately with EcoRI and HindIII at 37°C for 2 hours under standard conditions. The resulting four fragments of approximately 2.5 kb were isolated separately after gel electrophoresis by electroelution. The fragments were extracted with phenol and chloroform, and precipitated with ethanol. The four fragments were then resuspended in 10µl TE.

#### B. Ligation of expression cassettes into Bin19

Aliquots containing the four fragments from (A) were ligated to Bin19 DNA prepared as described in Example 15 (B) in separate ligation reactions under standard conditions. The ligation mixtures were used to transform competent TG2 cells. The transformation mixture was plated onto L-agar plates containing kanamycin and Xga1. After overnight incubation recombinant colonies were identified by their white colour. A number of clones for the separate experiments were picked and DNA was prepared. The DNAs were analysed for the presence of the appropriate EcoRI - HindIII fragments.

### EXAMPLE 18

Construction of PG vectors pJR76A and pJR76S

## A. Isolation of fragment (c)

10 ug gTOM 23 (a genomic clone containing the PG gene, NCIB No 12373) was cut with HindIII and BamHI. The 1.98 Kb fragment was isolated after agarose gel electrophoresis by electroelution. The  
 5 cohesive ends of the fragment were filled in with T<sub>4</sub> DNA polymerase.

## B. Ligation of fragment (c) to pJR1

10 The products from (A) above and Example 14 (B) (ie. pJR1 cut with SmaI) were ligated at 16 C for 24 hours under standard conditions and the mixture used to transform competent TG2 cells which were then plated onto plates containing ampicillin. Transformed colonies were grown up and used to prepare plasmid DNA. The DNA was cut with EcoRI and the orientation of the insert determined from the pattern of  
 15 fragments obtained. The clones were called pJR76A and pJR76S according to the orientation of the insert.

## EXAMPLE 19

Transfer of vectors pJR76A and pJR76S to Bin19

20

## A. Preparation of expression cassettes from pJR76A and pJR76S

5 ug of each clone was cut with HindIII at 37°C for 2 hours under standard conditions. The enzyme was inactivated by heating the reaction mixture to 70°C for 15 minutes. EcoRI was then added in concentration  
 25 necessary to give partial restriction. The reactions were stopped by the addition of phenol and chloroform. The required 2.71 Kb EcoRI - HindIII fragments were isolated after agarose gel electrophoresis by electroelution. The fragments were extracted with phenol and chloroform and precipitated with ethanol. The fragments were then resuspended in 10 ul TE.

30

## B. Ligation of the expression cassettes to Bin19.

The two fragments from vectors pJR76A and pJR76S prepared in (A) were ligated separately to Bin19 (prepared as described in Example 11 B). The ligation mixture was used to transform competent TG2 cells.  
 35 The transformation mix was plated onto L plates containing kanamycin and Xga1. Recombinant plasmids were identified by their white colour. DNA was prepared from a number of these and analysed for the presence of the required EcoRI - HindIII fragments.

## 40 EXAMPLE 20

Construction of PG vectors pJR86A and pJR86S

## A. Ligation of PG fragment (c) to pJR2

45

The products of Example 18 (A) (fragment c) and Example 16 (A) (ie pJR2 cut with HincII) were ligated at 16°C for 24 hours under standard conditions and the mixture used to transform competent TG2 cells which were then plated on plates containing ampicillin. Transformed colonies were grown up and used to prepare plasmid DNA. The orientation of the insert was deduced using the EcoRI restriction pattern. These  
 50 clones were called pJR86A and pJR86S, according to the orientation of the insert.

## EXAMPLE 21

55

Transfer of vectors pJR86A and pJR86S into Bin19.

This Example was carried out essentially as described in Example 19, ie. the vectors were cut with HindIII under conditions of partial restriction, which was then followed by restriction with EcoR1. The resulting 3.63 Kb fragment was isolated and cloned into Bin19.

All constructs in Bin19 were intended for use in separate triparental mating experiments to allow transfer to Agrobacterium, and from there to tomato plants.

## INHIBITION OF PECTIN ESTERASE

In addition to polygalacturonase, pectin esterase (PE) has been implicated in softening of the tomato fruit. A ripe tomato fruit cDNA library was screened with mixed oligonucleotide probes designed from the published amino acid sequence of PE. One clone, pPE1, (NCIB Accession No 12568) has been isolated and characterised. Figure 3 shows the complete sequence of this cDNA clone. The deduced amino acid sequence of PE is substantially different from the sequence published by Markovic and Jornvall. 40 amino acid differences are found in the sequence of the mature PE protein; major rearrangements in the continuity of the published amino acid sequence are evident and an additional 11 amino acids are found in the protein presented here. In addition to the sequences of the mature PE, both N-terminal and C-terminal amino acid extensions are detected in the polypeptide encoded by pPE1.

We have used fragments of the cDNA to construct antisense and sense vectors. These are summarised in Table 2.

Table 2

Vectors based on	Name of anti- sense vector	Name of sense vector	Fragment
pJR1	pJR101A	pJR101S	420 bp PstI
	pJR111A	pJR111S	351 bp BbvI
pJR2	pJR102A	pJR102S	420 bp PstI
	pJR112A	pJR112S	351 bp BbvI

## CONSTRUCTION OF ANTISENSE PE VECTORS

### EXAMPLE 30

#### Preparation of pJR101A and pJR101S

##### A. Isolation of a 420 bp fragment from pPE1

Plasmid pPE1 was cut with PstI at 37°C for 2 hours under standard conditions. The 420 bp PstI fragment was isolated after agarose gel electrophoresis by electroelution, extracted with phenol and chloroform and precipitated with ethanol. The DNA was then resuspended in 10µl TE.

## B. Preparation of pJR1

pJR1 (from Experiment 11) was cut with PstI at 37°C for 2 hours under standard conditions. The reaction was stopped by the addition of phenol, precipitated with ethanol and resuspended in 20µl TE.

## C. Ligation of PE fragment to pJR1

The products of steps (A) and (B) above were ligated under standard conditions and the ligation mixture was used to transform competent TG2 cells. The transformation mix was subsequently plated onto ampicillin containing plates and incubated at 37°C overnight. Transformed colonies were grown up and used to prepare plasmid DNA. Clones were identified which gave 420bp fragment on digestion with PstI. The 650 bp BamHI - HindIII fragments from these clones were isolated after agarose gele electrophoresis by electroelution and cloned into M13mp8. The orientation of the PstI insert was determined by sequence analysis. Clones identified were named pJR101A and pJR101S according to the orientation of the insert.

## EXAMPLE 31

### 20 Transfer of vector pJR101A and pJR101S to Bin19

#### A. Isolation of a 1.2 Kb EcoR1 - HindIII fragment

Plasmids pJR101A and pJR101S were cut separately with EcoR1 and HindIII at 37°C for 2 hours under standard conditions. The resulting 1.2 Kb fragments were isolated after gel electrophoresis from agarose gels by electroelution. The DNA was then extracted with phenol and chloroform, precipitated with ethanol, and resuspended in 20 ul TE.

#### 30 B. Preparation of Bin19

Bin19 was cut with EcoR1 and HindIII for 2 hours at 37°C under standard conditions. The enzymes were removed by phenol extraction and the vector precipitated with ethanol. The DNA was then resuspended in water.

#### C. Ligation of the PE expression cassette to Bin19

Aliquots of the products of reactions A and B were ligated for 16 hours at 16°C under standard conditions. The ligation mix was used to transform competent TG2 cells. The transformation mix was plated onto plates containing kanamycin. DNA was picked from individual clones and analysed for the presence of the 1.2 Kb EcoR1 - HindIII fragment.

## 45 EXAMPLE 32

### Construction of vectors pJR102A and pJR102S

The construction of these vectors followed the construction of pJR101A and pJR101S (Example 30), except that the 420 bp PstI PE fragment was inserted into pJR2 (from Experiment 14).

## EXAMPLE 33

### 55 Transfer of vectors pJR102A and pJR102S to Bin19

Transfer of the PE expression cassettes was carried out as described in Example 31 for the transfer of pJR102A and pJR102S into Bin19.



## EXAMPLE 34

## Construction of vectors pJR111A and pJR111S

## 5 A. Isolation of a 351 bp fragment from pPE1.

Plasmid pPE1 was cut with BbvI at 37°C for 2 hours under standard conditions and the cohesive ends filled using T4 polymerase. The 351 bp fragment was isolated after agarose gel electrophoresis by electroelution, extracted with phenol and chloroform and precipitated with ethanol. It was then resuspended  
10 in 10 µl TE.

## B. Preparation of pJR1

15 1 µg pJR1 (from Experiment 11) was cut with SmaI for 2 hours at 37°C under standard conditions. The reaction was terminated by the addition of phenol and chloroform. After 5 extraction the DNA was precipitated with ethanol, and resuspended in 10 µl TE.

## 20 C. Ligation of the PE fragment to pJR1

The products of (A) and (B) were ligated at 16°C for 24 hours under standard conditions. The ligation mix was used to transform competent E.coli TG2 cells. The transformation mix was plated onto ampicillin-containing plates. Single colonies were grown up and analysed for the presence of a 900bp EcoRI - PstI  
25 fragment. This fragment was isolated by electroelution after agarose gel electrophoresis and cloned into M13 mp8 (commercially available vector). The orientation of the fragment was determined by DNA sequence analysis.

## 30 EXAMPLE 35

## Transfer of vectors pJR111A and pJR111S to Bin 19

## A. Isolation of the 1.1Kb EcoRI - HindIII fragment

35 Plasmids pJR111A and pJR111S were cut with EcoRI and HindIII at 37°C for 2 hours under standard conditions. The 1.1 Kb fragment was isolated after agarose gel electrophoresis by electroelution. It was extracted with phenol and chloroform, precipitated with ethanol and resuspended in 20 µl TE.

40 B. Ligation of the PE expression cassettes into Bin19.

Aliquots of the products of (A) and Example 31 (B) were ligated at 16°C for 24 hours. The ligation mixtures were used to transform competent E.coli TG2 cells. The transformation mix was plated onto plates  
45 containing kanamycin. Single colonies were used for DNA extraction and clones identified by the presence of the 1.1 Kb EcoRI - HindIII fragment.

## EXAMPLE 36

50 Construction of vectors pJR112A and pJR112S

Construction of these vectors followed the procedure in Example 34 for the construction of vectors pJR111A and pJR111S except that the 351 bp BbvI fragment was inserted into pJR2 from Experiment 14,  
55 rather than pJR1.


## EXAMPLE 37

## FIG. 1.

## DNA sequence of pTom 6

AATCTTTTCAATAGACAAGTTTAAAAACCATACATATAACAATATATCATGTTATCCAAAGGAATAGTATTCTCCTTCTCATTATTATTTTCTTCATCAATTTCAACTTGTAGAA  
 10 20 30 40 50 60 70 80 90 100 110 120  
 NVIDDNLFLKQVYDNILEQEFAHDFQAYLSYLSKNIESNN  
 GCAATGTTATTGATGACAATTTATCAACAAGTTTATGATAATTTCTTGAACAAGATTTGCTCATGATTTTCAAGCTTATCTTTCTTATTGAGCAAAAATATTGAAGCAACATA  
 130 140 150 160 170 180 190 200 210 220 230 240  
 IDKVDKNGIKVINVLFSFBAKGDGKTYDNIAFEQAHNEAC'S  
 ATATTGACAAGTTGATAAAATGGGATTAAAGTGAATGACTTAGCTTTGGAGCTAAGGGTGAAGGAAAAATATGATAATTTGATTTGAGCAAGCATGGAATGAAGCATGTT  
 250 260 270 280 290 300 310 320 330 340 350 360  
 SRTPVQFVUPKKNKYNLLKQITFSBPCRSSISVKIFGSLEA  
 CATCTAGAACCTGTTCAATTTGTGTTCTTAAAAACAAGATTATCTTCAAGCAATCACCTTTTCAGGTCATGCAATCTTCTATTTCAGTAAAGATTTTGGATCCTTAGAAG  
 370 380 390 400 410 420 430 440 450 460 470 480  
 SSKISDYKDRRLWIAFDSVQNLVUVGGGGTINGNGQVWHPS  
 CATCTAGTAAATTTGAGACTACAAAGATAGAAGGCTTTGGATTGCTTTTGATAGTGTTCAAATTTAGTTGTTGGAGGAGGAGAACTATCAATGGCAATGGACAAGTATGGTGGCAA  
 490 500 510 520 530 540 550 560 570 580 590 600  
 SCKINKSLPCRDAPALTFFHNCKNLKVNNLKS KNAQQIHI  
 GTTCTTGCAAAATAAATAATCACTGCCATGCAAGGATGACCAACGGCTTAACCTTCTGGAATTCGCAAAATTTGAAAGTGAATATCTAAGAGTAAAAATGCACACCAATTCATA  
 610 620 630 640 650 660 670 680 690 700 710 720  
 KFFSCTNVUASNLMINASAKSPNTDGVHVVSNTQYIQISDT  
 TCAATTTGAGTCATGCAATATGTTGATCTCAATTTGATGATCAATGCTTCAGCAAGAGGCCAATACTGATGGAGTCCATGATCAATCTCAATATTCATATCTGATA  
 730 740 750 760 770 780 790 800 810 820 830 840  
 IIGTGDDCISIVSBSQNVQATNITCGBPBGISIGSLGSGN  
 CTATTATTGGAAGGTGATGATTGATTTCAATTTGTTCTGGATCTCAAAATGTGAGGCCACAAATATTACTTGTGTTCCAGGTCATGTTAGTATTGGAAGCTTAGGATCTGGA  
 850 860 870 880 890 900 910 920 930 940 950 960  
 SEAYVSNVTUNEAKIIGAENBVRIKTHWQGBSGQASNIKFL  
 ATTCAGAGCTTATGTCTAATGTTACTGTAATGAAGCCAAATTTATCGGTGCCGAAATGAGTTAGGATCAAGCTTGGCAGGAGGATCTGGACAAGCTAGCAACATCAATTTTC  
 970 980 990 1000 1010 1020 1030 1040 1050 1060 1070 1080  
 NVEHQDVKYPIIIDQNYCDRVEPCIQQFS AVUKNVUYEN  
 TGAATGTGGAATGCAAGAGCTTAAGTATCCATAATTATAGCAAACTATTGTGATCGAGTTGAACATGTATACACAGTTTTCAGCAATTCAGTGAAGTGAAGTGTGATGAGA  
 1090 1100 1110 1120 1130 1140 1150 1160 1170 1180 1190 1200  
 IKSTSA TKVAIKFDCSTNFPCEGIIMENINLVGESGKPS E  
 ATATCAAGGBCAAGTGCAACAAAGGTGCCATAAAATTTGATTGCAACCAAACTTTCCATGTGAAGGAATTATAATGAGAAATATAATTTAGTAGGGGAAAGTGGAAAAACCATCAG  
 1210 1220 1230 1240 1250 1260 1270 1280 1290 1300 1310 1320  
 ATCKNVHFNNAEHVTPHCTSLSEI SEDEALLYNY\*  
 AGGCTACGTGCAAAATGTCCATTTTAACAATGCTGAACATGTTACACCACACTGCATTCCTAGAAATTTGAGAGGATGAAGCTCTTTTGTATAATTATTAATTATATCTATAGATCT  
 1330 1340 1350 1360 1370 1380 1390 1400 1410 1420 1430 1440  
 TCAATATATAGCAGATATGATATATCACAATAAACAATCTATATCTATGATTGAATAATTATTATTAATATGATGAGGATTGAAGTTTTAATAGACTACTATGATTTTCTATTTTCTA  
 1450 1460 1470 1480 1490 1500 1510 1520 1530 1540 1550 1560  
 GTCAAAAGTTTGACGATTGACTTTTTATGTACAAAAATAAATAAATGTTTATTATATGAAAAAAAAAAAA  
 1570 1580 1590 1600 1610 1620 1630

### PG gene and cDNA fragments used in the construction of PG antisense vectors



## FIG. 3.

## DNA sequence of pPE1

CTTTAACTGATTGTCTTGAGCTTCTTGATCTGTGAGTTGATTTAGTATGTGATTCAATTGCAGCAATTGATAAGAGAAGTCSTTCGGAGCATGCCAATGCCAAAGTTGGCTAAGTGGT  
 10 20 30 40 50 60 70 80 90 100 110 120  
 TGCTTACTAACCACGTTACGTGCTTGGATGAGCTTGATTCTTTACTAAGCTATGATAAATGGACCAATCTTGATGAGTTGATCTCGAGAGCTAAGGTAGCATTGGCGATGCTTGCCT  
 130 140 150 160 170 180 190 200 210 220 230 240  
 V T T P N D E V L R P G L G K M P S W U S S R D R K L M E S S G K D I G A N A V  
 CTGTGACAACTCCAAATGATGAAGTTTGAAGCCGGGTTTAGGAAAATGCCATCTTGGGTGAGTTGAGGGATAGGAAGCTGATGGAGAGTTGCGGTAAAGGACATTGGAGCGAATGCAG  
 250 260 270 280 290 300 310 320 330 340 350 360  
 V A K D G T G K Y R T L A E A V A A P D K S K T R Y V I Y V K R B T Y K E N V  
 TGGTGGCAAAAGATGGAACAGGGAATATCGAACCTTGTCTGAAGCTGTGCTGCAGCACCAGATAAGAGTAAGACGCGTTATGTAATTTATGTAAGAGGGGAAGCTTATAAGAGAGATG  
 370 380 390 400 410 420 430 440 450 460 470 480  
 E V S S R K M N L M I I G D G H Y A T I I T G S L N V U D B S T T F H S A T L A  
 TTGAGGTGAGTAGCAGGAAATGATTTGATGATTATTGGTGTGATGGCATGTATGCTACCATCATTTACTGGAGGCTTAATGTTGTGATGGATCAACAACCTTCCACTCTGCCACTCTTG  
 490 500 510 520 530 540 550 560 570 580 590 600  
 A V G K G F I L Q D I C I Q N T A B P A K H Q A V A L R V B A D K S V I N R C R  
 CTGCAGTTGGCAAGGATTTATACTACAGGACATATGTATACAGAACACAGCAGGACGAGCTAAGACCAAGCTGTTGCAGTTGAGGTGAGAGCTGATGAAGCTGTGCATAAATCGTTGTC  
 610 620 630 640 650 660 670 680 690 700 710 720  
 I D A Y Q D T L Y A H S Q R Q F Y Q S S Y V T G T I D F I F G N A A V V F Q K C  
 GTATCGATGCTTATCAAGACCCCTTTATGACATTCTCAAGGCAATTCATCAGAGCTCCTACGTGACAGGGCAATTGATTTCATATTGCGTAATGCAGCAGTTGATTTCCAGAAAT  
 730 740 750 760 770 780 790 800 810 820 830 840  
 Q L V A R K P G K Y Q Q N M V I A Q G R T D P N Q A T G T S I Q F C D I I A S P  
 GCCAGCTGTAGCTAGAAAACCGGGTAATACAGCAAAACATGCTGATCGACAAGGACGAGGACCAATCAGGCCACGGGACATCAATTGATTTTGTGATATAATAGCAAGTC  
 850 860 870 880 890 900 910 920 930 940 950 960  
 D L K P V V K E F P T Y L G R P H K K Y S R T V V H E S S L B G L I D P S B W A  
 CTGACCTAAACAGTCGTGAAGAATTCCCAACATATCTTGTAGGCCATGGAAAAATTAAGAACTGTAGTGAATCATCATTGGGTGCTCATTGATCCATCGGGTTGGG  
 970 980 990 1000 1010 1020 1030 1040 1050 1060 1070 1080  
 E H H G D F A L K T L Y Y G E F M N N G P B A G T S K R V K H P G Y H V I T D P  
 CTGAGTGGCAGGAGATTTTGTGTTAAGACATTGATTATGGTGAATTTATGAATAATGACCTGGTGTCTGCTAGTAAAGCTGTCAAGTGGCTGGCTATCATGTCTACTGACC  
 1090 1100 1110 1120 1130 1140 1150 1160 1170 1180 1190 1200  
 A E A H S F T V A K L I Q G G S H L R S T D V A Y V D B L Y D Y S D I K L L F V  
 CCCTGAAGCTATGTCTTCACTGTGGCTAAGCTGATTGAGGCGGATCATGGTTGAGGTCTACTGACGTGGCSTATGTGGATGGATTATATGATTATAGTATATAAATTACTCTTTG  
 1210 1220 1230 1240 1250 1260 1270 1280 1290 1300 1310 1320  
 Y V T R H L \*  
 TTTATGTAACAGACATCTTAAAAAGTTCAAGTAAGTAGTAGTAATATATCATATGAAAGTCCACATGAGCAGGGCAGAGCGGATTAAGTGTCTAAGCATAACACACAACCTCTAGT  
 1330 1340 1350 1360 1370 1380 1390 1400 1410 1420 1430 1440  
 GTGACAAGCATTACATGGCTCATTCTTACTACTAAGTCGTCAATAAGTTCAAGTTAAGGGGTTCAAGTTAATATACGTATATATTTATGTTGGCGATAAAGCTGAGTGTATG  
 1450 1460 1470 1480 1490 1500 1510 1520 1530 1540 1550 1560  
 CTTTAACTGATTGTCTTGAGCTTCTTGATCTGTGAGTTGATTTAGTATGTGATTCAATTGCAGCAATTGATAAGAGAAGTCSTTCGGAGCATGCCAATGCCAAAGTTGGCTAAGTGGT  
 1570 1580 1590 1600 1610 1620 1630 1640 1650 1660 1670 1680

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(11) Publication number:

**0 271 988  
A3**

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: 87309853.7

(51) Int. Cl.4: **C12N 15/00 , C12N 5/00 ,  
A01H 1/00**

(22) Date of filing: 06.11.87

(30) Priority: 11.11.86 GB 8626879

(43) Date of publication of application:  
22.06.88 Bulletin 88/25

(84) Designated Contracting States:  
AT BE CH DE ES FR GB GR IT LI LU NL SE

(88) Date of deferred publication of the search report:  
26.04.89 Bulletin 89/17

(71) Applicant: **IMPERIAL CHEMICAL INDUSTRIES  
PLC**

Imperial Chemical House Millbank  
London SW1P 3JF(GB)

(72) Inventor: **Bridges, Ian George  
Greentrees**

Manley Lane Cheshire(GB)  
Inventor: **Schuch, Wolfgang Walter**  
20 Belmonte Road Overton  
Frodsham Cheshire(GB)

Inventor: **Grierson, Donald**  
6 Tyler Court  
Shepshed Leicestershire(GB)

(74) Representative: **Roberts, Timothy Wace et al**  
Imperial Chemical Industries PLC Legal  
Department : Patents PO Box 6 Bessemer  
Road  
Welwyn Garden City Herts, AL7 1HD(GB)

(54) **Anti-sense regulation of plant gene expression.**

(57) Recombinant DNA comprises promoter and terminator base sequences respectively upstream and downstream of an inverted base sequence complementary to a substantial sequence of bases in polygalacturonase mRNA. The antisense mRNA produced thereby delays softening of fruit, in particular tomatoes.

EP 0 271 988 A3



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number

EP 87 30 9853

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
D, Y	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE USA, vol. 83, no. 15, August 1986, pages 5372-5376, Washington, US; J.R. ECKER et al.: "Inhibition of gene expression in plant cells by expression of antisense RNA" * Whole article *	1-12	C 12 N 15/00 C 12 N 5/00 A 01 H 1/00
Y	NATURE, vol. 315, 13th June 1985, pages 601-603; J. COLEMAN et al.: "A novel immune system against bacteriophage infection using complementary RNA (micRNA)" * Page 306, right-hand column, lines 5-12 *	1-12	
O, Y	CHEMICAL ABSTRACTS, vol. 107, 1987, page 388, no. 74243f, Columbus, Ohio, US; D. GRIERSON et al.: "Expression and function of ripening genes", & PLANT BIOL. 1987, 4(TOMATO BIOTECHNOL.), 309-23 * Abstract *	1-12	
Y	BIOLOGICAL ABSTRACTS/RRM, no. 29054425; W. SCHUCH et al.: "Isolation and characterization of complementary DNA and genomic clones for tomato polygalacturonase", & JOURNAL OF CELLULAR BIOCHEMISTRY SUPPLEMENT(US), 1985, vol. 0, no. 9, part C, page 241 * Abstract *	8	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
			C 12 N A 01 H
Place of search THE HAGUE		Date of completion of the search 20-01-1989	Examiner MADDOX A. D.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
A	CHEMICAL ABSTRACTS, vol. 106, 1987, no. 81504z, Columbus, Ohio, US; D. GRIERSON et al.: "Gene expression during tomato ripening", & PHILOS. TRANS. R. SOC. LONDON B 1986, 314(1166), 399-410 * Abstract * ---	2-6	
A	CHEMICAL ABSTRACTS, vol. 93, 1980, page 375, no. 64543g, Columbus, Ohio, US; R.B. WILLS et al.: "Effect of calcium on activity of mitochondria and pectic enzymes isolated from tomato fruits", & J. FOOD BIOCHEM. 1980, 3(2-3), 103-10 * Abstract * ---	2	
A	CHEMICAL ABSTRACTS, vol. 101, 1984, page 416, no. 127096v, Columbus, Ohio, US; G.E. HOBSON et al.: "The inhibition of tomato fruit ripening by silver", & J. PLANT PHYSIOL. 1984, 116(1), 21-9 * Abstract * ---	2	
A	CHEMICAL ABSTRACTS, vol. 95, 1981, page 364, no. 184019d, Columbus, Ohio, US; D. GRIERSON et al.: "The regulation of gene expression during the ripening of tomato fruits", & LONG ASHTON SYMP. [PROC.] 1981, 7th(QUAL. STORED PROCESSED VEG. FRUIT), 179-91 * Abstract * --- -/-	2	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
Place of search THE HAGUE		Date of completion of the search 20-01-1989	Examiner MADDOX A.D.
<b>CATEGORY OF CITED DOCUMENTS</b> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
A	CHEMICAL ABSTRACTS, vol. 97, 1982, page 321, no. 36188j, Columbus, Ohio, US; G.A. TUCKER et al.: "Purification and changes in activities of tomato pectinesterase isoenzymes", & J. SCI. FOOD AGRIC. 1982, 33(4), 396-400 * Abstract *	2	
A	CHEMICAL ABSTRACTS, vol. 104, 1986, no. 15979r, Columbus, Ohio, US; A. SLATER et al.: "Isolation and characterization of cDNA clones for tomato polygalacturonase and other ripening-related proteins", & PLANT MOL. BIOL. 1985, 5(3), 137 47 * Abstract *	1-6	
P,A	NUCLEIC ACIDS RESEARCH, vol. 14, no. 21, 11th November 1986, pages 8595-8603; D. GRIERSON et al.: "Sequencing and identification of a cDNA clone for tomato polygalacturonase" * Whole article *	1-6	
P,X	EP-A-0 240 208 (CALGENE) * Example 2; claims 4,5 *	1-6,9,10	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
Place of search THE HAGUE		Date of completion of the search 20-01-1989	Examiner MADDOX A.D.
<b>CATEGORY OF CITED DOCUMENTS</b> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			